

Sect. *c.* *Lamina of lateral wing broad, marginal ridge feebly developed; second to eleventh dorsal segments multisulcate.*

23. *ZEPHRONIA SULCATULA.*

*Zephronia sulcatula*, Butler, Ann. & Mag. Nat. Hist. ser. 4, vol. x. p. 357. n. 13, pl. 18. f. 5 (1872).

Borneo (*W. Jeakes*).

Type, B.M.

EXPLANATION OF PLATE XIX.

Fig. 1. *Sphærotherium glabrum*, profile.

1 a. — — —, anterior segments from above.

2. — — — *fraternum*, profile.

3. — — — *stigmaticum*, perspective view from behind.

4. *Zephronia noticeps*, head, nuchal plate, and anterior segments, enlarged; front view.

5. — — — *lutescens*, profile.

6. — — — *ignobilis*, profile, enlarged.

7. — — — *pilifera*, profile.

8. — — — *corrugata*, profile.

9. — — — *banksiana*, profile.

10. — — — *leopardina*, profile.

5. Descriptions of Eight new Species of Land and Marine Shells from various Localities. By GEORGE FRENCH ANGAS, Corr. Mem. Z.S., F.R.G.S., F.L.S., &c.

[Received December 16, 1872.]

(Plate XX.)

1. *EUTHRIA ARACANENSIS*, n. sp. (Plate XX. fig. 1.)

Shell ovately fusiform, solid, transversely ridged, the interstices finely transversely striate, the ridges stronger in front, pale olivaceous brown, ornamented with longitudinal irregular chestnut markings; whorls 9 to 10, strongly angulate and nodose in the middle, excavated above, and slightly rounded below; aperture ovate, nearly half the length of the shell, angulate posteriorly, pale flesh-colour; outer lip crenulate, finely sulcate within; columella strongly arcuate, with a moderate posterior callus; the rostrum slanting, recurved.

Length 2 inches, diam. 1 inch (*coll. Hanley*).

*Hab.* Aracan.

2. *HELIX (SEMICORNU) SILENUS*, n. sp. (Plate XX. fig. 2.)

Shell deeply and perspectively umbilicated, rather thin, globosely depressed, pale brown, minutely and very finely punctately striate; spire immersed; whorls 5, rounded, rapidly increasing, the last inflated, a little flattened at the upper part, descending considerably in front, and somewhat angulated round the umbilicus; aperture oblique, rhomboidly lunate; peristome thin, whitish; margins approximating, the right flexuous, narrowly expanded; the columellar margin rounded, expanded, and slightly reflexed.

Maj. diam. 7, min. 6; height 5 lines.

*Hab.* New Ireland.



Sowerby lith

M & N Hanhart. imp.

